

Meningococcal Disease and Camp Attendees: Commonly Asked Questions

What is meningococcal disease?

Meningococcal disease is caused by infection with bacteria called *Neisseria meningitidis*. These bacteria can infect the tissue (the "meninges") that surrounds the brain and spinal cord and cause meningitis, or they may infect the blood or other organs of the body. Symptoms of meningococcal disease can include fever, severe and constant headache, stiff neck or neck pain, nausea and vomiting, and rash. In the US, about 350-550 people get meningococcal disease each year and 10-15% die despite receiving antibiotic treatment. Of those who survive, about 10-20% may lose limbs, become hard of hearing or deaf, have problems with their nervous system, including long term neurologic problems, or have seizures or strokes.

How is meningococcal disease spread?

These bacteria are passed from person-to-person through saliva (spit). You must be in close contact with an infected person's saliva in order for the bacteria to spread. Close contact includes activities such as kissing, sharing water bottles, sharing eating/drinking utensils or sharing cigarettes with someone who is infected; or being within 3-6 feet of someone who is infected and is coughing and sneezing.

Who is most at risk for getting meningococcal disease?

People who travel to certain parts of the world where the disease is very common, microbiologists, people with HIV infection and those exposed to meningococcal disease during an outbreak are at risk for meningococcal disease. Children and adults with damaged or removed spleens or persistent complement component deficiency (an inherited immune disorder) are at risk. Adolescents, and people who live in certain settings such as college freshmen living in dormitories and military recruits are at greater risk of disease from some of the serotypes.

Are camp attendees at increased risk for meningococcal disease?

Children attending day or residential camps are **not** considered to be at an increased risk for meningococcal disease because of their participation.

Is there a vaccine against meningococcal disease?

Yes, there are 2 different meningococcal vaccines. Quadrivalent meningococcal conjugate vaccine (Menactra and Menveo) protects against 4 serotypes (A, C, W and Y) of meningococcal disease. Meningococcal serogroup B vaccine (Bexsero and Trumenba) protects against serogroup B meningococcal disease, for age 10 and older.

Should my child or adolescent receive meningococcal vaccine?

That depends. Meningococcal conjugate vaccine (Menactra and Menveo) is routinely recommended at age 11-12 years with a booster at age 16. In addition, this vaccine may be recommended for children with certain high-risk health conditions, such as those described above. Otherwise, meningococcal vaccine is **not** recommended for attendance at camps.

Meningococcal serogroup B vaccine (Bexsero and Trumenba) is recommended for people with certain relatively rare high-risk health conditions (examples: persons with a damaged spleen or whose spleen has been removed, those with persistent complement component deficiency (an inherited disorder), and people who may have been exposed during an outbreak). Adolescents and young adults (16 through 23 years of age) who do not have high risk conditions may be vaccinated with a serogroup B meningococcal vaccine, preferably at 16 through 18 years of age, to provide short term protection for most strains of serogroup B meningococcal disease. Parents of adolescents and children who are at higher risk of infection, because of certain medical conditions or other circumstances, should discuss vaccination with their child's healthcare provider.

How can I protect my child or adolescent from getting meningococcal disease?

The best protection against meningococcal disease and many other infectious diseases is thorough and frequent handwashing, respiratory hygiene and cough etiquette. Individuals should:

1. wash their hands often, especially after using the toilet and before eating or preparing food (hands should be washed with soap and water or an alcohol-based hand gel or rub may be used if hands are not visibly dirty);
2. cover their nose and mouth with a tissue when coughing or sneezing and discard the tissue in a trash can; or if they don't have a tissue, cough or sneeze into their upper sleeve.
3. not share food, drinks or eating utensils with other people, especially if they are ill.
4. contact their healthcare provider immediately if they have symptoms of meningitis.

If your child is exposed to someone with meningococcal disease, antibiotics may be recommended to keep your child from getting sick.