

6-Step Return to Play Progression

It is important for an athlete's parent(s) and coach(es) to watch for concussion symptoms after each day's return to play progression activity. An athlete should only move to the next step if they do not have any new symptoms at the current step. If an athlete's symptoms come back or if he or she gets new symptoms, this is a sign that the athlete is pushing too hard. The athlete should stop these activities and the athlete's medical provider should be contacted. After more rest and no concussion symptoms, the athlete can start at the previous step.

Step 1: Back to regular activities (such as school)

Athlete is back to their regular activities (such as school).

Step 2: Light aerobic activity

Begin with light aerobic exercise only to increase an athlete's heart rate. This means about 5 to 10 minutes on an exercise bike, walking, or light jogging. No weight lifting at this point.

Step 3: Moderate activity

Continue with activities to increase an athlete's heart rate with body or head movement. This includes moderate jogging, brief running, moderate-intensity stationary biking, moderate-intensity weightlifting (less time and/or less weight from their typical routine).

Step 4: Heavy, non-contact activity

Add heavy non-contact physical activity, such as sprinting/running, high-intensity stationary biking, regular weightlifting routine, non-contact sport-specific drills (in 3 planes of movement).

Step 5: Practice & full contact

Young athlete may return to practice and full contact (if appropriate for the sport) in controlled practice.

Step 6: Competition

Young athlete may return to competition.